

GEOMETRIC PROOFS AND ALGEBRAIC SETS

DAVIDE MARAN

ABSTRACT. This article shows a new method to prove theorems inspired and based on some algebraic geometry results like the Nullstellensatz. This method allows to assert the validity of a geometry theorem by proving it only in some particular cases.

1. INTRODUCTION

In order to understand the proofs of this article, it is necessary to clarify some intuitive definitions about the two-dimensional cartesian coordinate system. Consider an *element* a defined set of points in the cartesian coordinate system, which means every object of the euclidean geometry except for angles.

Definition 1.1. Given an element α in Cartesian coordinate system we say that it *depends on* a set A of variables $a, b, c, \dots$ if that set identifies the element completely.

The condition "identifies completely" means that there must be a function f which links every ordered t -uple of the values of the variables to one and only one element described and this function must be surjective. However f does not have to be injective (and so bijective) as it is shown after the second example.

Because of the fact that we are considering the cartesian coordinate system, the variables must belong to the real field. This fact will cause some issues later.

Example 1.2. "the straight line r ", cannot always be written as $y = kx + q$ (this equation does not include the parallels to the y -axis), so we cannot say that it depends on $A\{k, q\}$.

Example 1.3. "the straight line r not parallel to the y -axis", as it can be written as $y = kx + q$, depends on $A\{k, q\}$.

2010 *Mathematics Subject Classification.* Primary 51N20; Secondary 03F07.

Key words and phrases. Geometry, Proof, Polynomial.

Note that it is already well known that any element can be represented using different sets and different types of variables. For example the straight line r can also depend on the set $B\{a, b, c, d\}$ where $P(a, b)Q(c, d)$ are two points of r .

Proposition 1.4. *for every triangle EFG , there exists at least one ordered triple of a, b, c so that ABC $A(0, 0)B(b, a)C(c, 0)$ is congruent to EFG*

Proof. Move E to the Origin with a traslation, and rotate EFG until EG belongs to the axis x . Then let $c = |EG|$. now the point F is fixed in one point of the plane and let (u, v) be its coordinates. Let $b = u, a = v$ then $E(0, 0), F(b, a), G(c, 0)$, so it is congruent to ABC . $\square$

Due to this fact it is possible to add one more fundamental example which will result extremely impotant.

Example 1.5. The generic triangle, as it can be written as ABC $A(0, 0)B(b, a)C(c, 0)$ depends on $\{a, b, c\}$.

Definition 1.6. Given a predicate E , if all the elements of E depend on a set of variables $A\{a, b, c\}, \dots$ Or on one of its subsets, we say that the predicate *depends on* the set of variables $A\{a, b, c, \dots\}$

According to this perspective, we interpret every predicate as a relation between geometric constructions made starting from a set of variables.

Example 1.7. "two points A, B and the Origin are collinear", if we put $A(a, b)B(c, d)$, depends on $A\{a, b, c, d\}$.

Definition 1.8. Given a predicate in the cartesian coordinate system which depends on the variables $a, b, c, \dots$ we call *associated condition* (C.A.) the set of values that satisfy the statement.

As said above, as there are multiple ways to choose the variables, there are multiple different associated conditions with the same predicate but, for evry set of variables there is only one C.A. The associated condition can either be the set of solution of an equation or the set of solution of an inequality; in the first case this equation can be algebraic or trascendental. In practice, if we deal with a predicate of euclidean geometry

as the formulas used to "build" the elements of the problem (parallel lines, middle points, bisectors...) are reducible to polynomial equations, the condition associated can usually be reduced to a polynomial equation. This article does not consider the predicates which can be associated to an inequality, for example "the point A is nearer to point B than to the straight line r ". Before applying every theorem that follows it is needed to verify that the statement has an equation as associated condition.

Example 1.9. "two points $A(a, b)$ and $B(c, d)$ and the origin are collinear" depends on $A\{a, b, c, d\}$.

If $a = c = 0$ E is verified.

Otherwise the statement is valid if and only if the straight lines r and s , passing for OA and OB respectively, have the same angular coefficient. This holds when $b/a = d/c$, so the associated condition is the set of solutions of:

$$bc = ad$$

However, despite there can be only one set of values which satisfy a predicate E, we cannot say that the polynomial equation that we have found is *the* equation associated to this statement, neither that it is the more simple. For example, the equation $(bc - ad)^2 = 0$, as it has exactly the same solutions, can represent the set as well.

In order to solve this problem we should specify that we call *associated polynomial* the polynomial with real coefficients and the less degree between all the ones which have the C.A. in their set of solutions. This means that the associated polynomial has the same set of real zeros of the one we found after the calculus required from the problem. This definition is well-posed because two polynomials of the same degree, with the same solutions, can be made equal by multiplication for a constant term, so there is only one associated polynomial. From this point on it would be possible to call associated condition also the equation which equals the polynomial associated to zero, because its solutions correspond exactly to C.A. intended as set of tuples.

Still, anyway, we are not able to determine this polynomial associated. In order to solve this problem it is needed a brief introduction to one of the most important results in the algebraic geometry.

2. CONSIDERATIONS AND RESULTS CONCERNING THE ALGEBRAIC GEOMETRY

Definition 2.1. Let $C[x_1, x_2, x_3, \dots, x_n]$ (C stands for the set of the complex numbers) be the ring of polynomials with coefficients in C and the variables $x_1, x_2, x_3, \dots, x_n$ in C .

Definition 2.2. Given a set S of polynomials in the ring, denote $V(S)$ the ordered tuples (points in the n -dimensions complex space) $P \in C^n$ such that

$$\forall f \in S, f(P) = 0$$

Definition 2.3. A subset $X \in C^n$ is called *algebraical set* if there exists a set S of polynomials in the ring so that $V(S) = X$

Definition 2.4. Let $I(U)$, where $U \in C^n$ the set S of polynomials in the ring such that

$$\forall f \in S, \forall u \in U, f(u) = 0$$

The I stands for ideal, because the most basilar result of the algebraic geometry is that the set $I(U)$ is an ideal of the ring $C[x_1, x_2, x_3, \dots, x_n]$. Since in the cartesian coordinate system the variables can only assume real values, we need to impose a condition under that the results of algebraic geometry can be applied in this case.

Definition 2.5. Let a *R-factorizable polynomial* (or polynomial equation) be a polynomial whose factorization in C contains only polynomials with real coefficients and has infinite solutions over R

Now the problem exposed at the end of the previous section can be better defined:

Call P the polynomial which can be found from the formulas of cartesian plane, $bc - ad$ in the example 1.9.

Let $S : \{P\}$ and let P be R -factorizable. The set of all the polynomials which include C.A. in their set of solutions is a subset of $I(V(S))$ because:

1) Under the assumption of R -factorizability $V(S)$ is the smallest algebraical set which includes the C.A. and in particular

$$\forall x_1, x_2, \dots, x_n \in R, (x_1, x_2, \dots, x_n) \in C.A. \iff P(x_1, x_2, \dots, x_n) = 0$$

2) $I(V(S))$ is the ideal which contains all the polynomials whose set of real solution includes C.A. ($V(S)$ contains some tuples of complex numbers which do not belong to C.A. however from the passage 1), we know that every polynomial Q which contains C.A. in its set of zeros, have to contain also the complex solutions).

The importance of this result is the fact that the form $I(V(S))$ allows us to apply one of the most important Hilbert's work (the proof will not be included as it digress from the aim of this article).

Theorem 2.6. (*Nullstellensatz*) *Let C be an algebraically closed field. Let $C[x_1, x_2, x_3, \dots, x_n]$ be the ring of polynomials with coefficients in C and $S \in C[x_1, x_2, x_3, \dots, x_n]$. Then*

$$\forall S, I(V(S)) = \sqrt{S}$$

Where $\sqrt{S}$ denote the radical of the ideal J generated by S , so the set $Z \in C[x_1, x_2, x_3, \dots, x_n]$ so that

$$\forall z \in Z, \exists n \in \mathbb{N} | z^n \in J$$

Now, since our polynomial associated is the "smallest" belonging to $I(V(S))$, we can find it in an easy way:

Proposition 2.7. *Let $S = \{P\}$. Let $[x_1, x_2, x_3, \dots, x_n] = \theta$, $P(x_1, x_2, x_3, \dots, x_n) = P(\theta)$. Let $P_1^n(\theta)P_2^m(\theta)\dots = P(\theta)$ be its factorization.*

Then $Q(\theta) = P_1(\theta)P_2(\theta)\dots$ is the polynomial with the lowest degree so that $Q(\theta) \in I(V(S))$

Proof. Let $E\{n, m, \dots\}$ be the set of the degrees of the unfactorizable polynomials in the factorization of P . Since E is finite set of integers, its maximum M exists and is still an integer. Then, thanks to the Nullstellensatz, since $Q(\theta)^M$ is multiple of P and so belongs to J , $Q(\theta) \in I(V(S))$. Furthermore since every polynomial with a lower degree is not multiple of one between $P_1(\theta), P_2(\theta)\dots$, there cannot be a Q_2 which satisfies the condition with a less degree. $\square$

Now we can answer the last question in the previous section:

Example 2.8. The polynomial associated with the predicate "two points $A(a, b)B(c, d)$ and the origin are collinear" is

$$bc - ad = 0$$

because this polynomial is not factorizable.

What is the importance of the definition of polynomial associated stays in this theorems:

Theorem 2.9. *Suppose that $P(x, y, z, \dots)$ and $G(x, y, z, \dots)$ are polynomials in the same variables, belonging to $C[x_1, x_2, x_3, \dots, x_n]$ so that for every $(x_0, y_0, z_0, \dots)$ if $P(x_0, y_0, z_0, \dots) = 0$ then also $G(x_0, y_0, z_0, \dots) = 0$. Let $P(x, y, z, \dots)$ so that his factorization contains only unfactorizable polynomials of degree one. Then $P(x, y, z, \dots)$ divides $G(x, y, z, \dots)$.*

Proof. Let $S = \{P\}$. The condition

$P(x_0, y_0, z_0, \dots) = 0 \Rightarrow G(x_0, y_0, z_0, \dots) = 0$ means that the set of solution of P is a subset of the set of solution of G . So $G(x, y, z, \dots) \in I(V(S))$. So, for the Nullstellensatz, there exists an $n \in \mathbb{N}$ so that $G^n(x, y, z, \dots)$ belongs to J (the ideal generated by P). This means, according to the definition of ideal of a ring generated by P , that $G^n(x, y, z, \dots)$ is multiple of $P(x, y, z, \dots)$ but, due to the Hypothesis about the factorization of P , this implies that

$G(x, y, z, \dots)$ is multiple of $P(x, y, z, \dots)$ □

Corollary 2.10. *Suppose E is a statement depending on the set of variables $\{a, b, c, \dots\}$ and let E_2 be another statement on the same variables so that $E \Rightarrow E_2$. Also let the condition associated with E_2 be written as $G(a, b, c, \dots) = 0$ (polynomial equation) and the condition associated with E be an R -factorizable polynomial equation $P(a, b, c, \dots) = 0$ where $P(a, b, c, \dots)$ is not dividable for any square polynomial. Then the polynomial associated with E divides the polynomial associated with E_2 .*

Proof. the two polynomials $G(a, b, c, \dots)$ and $P(a, b, c, \dots)$ respect all the conditions to apply theorem 2.9 □

Remark 2.11. In a real situation how can we reach the result that $E \Rightarrow E_2$?

Being E and E_2 predicates of geometry we have to do a geometrical proof of this fact. As the cartesian coordinate system describes the plane with variables which can only assume real values, then we have proved that

$$P(x, y, z, \dots) = 0 \Rightarrow G(x, y, z, \dots) = 0$$

(as above, $P(x, y, z, \dots) = 0$ is the condition associated with E and $G(x, y, z, \dots) = 0$ is the condition associated with E_2) only in the case

when $x, y, z, \dots \in R$. This is the reason why it is necessary to impose the condition of R-factorizability.

Example 2.12. This is what it would be possible without the condition of R-factorizability.

Let E be the predicate "Point $W(a, b)$ is the Origin".

Let E_2 be the predicate "Point $W(a, b)$ belongs to the line $y = x$ ".

Now, for every pair of $(a, b) \in R$ $a^2 + b^2 = 0 \Leftrightarrow E$. So $a^2 + b^2 = 0$ is the condition associated with E .

The condition associated with E_2 is $a - b = 0$.

Because of the fact that $E \Rightarrow E_2$ it is proven with corollary 2.10 that $a^2 + b^2$ divides $a - b$.

As theorem 2.9 imposes the condition "the polynomial is not dividable for any square polynomial", it is useful to introduce a new operation between polynomials that we will call m.c.s. which is a variant of L.C.M. whose result is a polynomial that is not dividable for any square polynomial. m.c.s. can be defined in terms of algebraic geometry as follows:

Definition 2.13. Let $A\{P(\theta), G(\theta), \dots\}$ be a subset of $C[x, y, z, \dots] = C[\theta]$. Then $m.c.s.(A)$ is the polynomial with the less degree which belongs to the radical of the ideal generated by A .

The operative definition of m.c.s. is:

Let $A\{C(x, y, z, \dots), D(x, y, z, \dots), \dots\}$.

- 1) Factorize L.C.M.(A) as product of irreducible polynomials.
- 2) Reduce all the factors to degree 1

Example 2.14. $L.C.M.[(x - y), (x + y)^2] = (x - y)(x + y)^2$

m.c.s. $[(x - y), (x + y)^2] = (x - y)(x + y)$

3. AUTARKY THEOREM

Here it is the main results of this article, which I call Autarky Theorem.

Main Theorem 3.1 (Autarky). *Let E be a predicate of geometry referred to the Cartesian coordinate system depending on the set A of variables*

$$S\{a, b, c, \dots\} = \theta$$

Call $E(a, b, c, \dots) = E(\theta)$ its associated polynomial. Let E_1, E_2, E_3 be predicates so that $\forall n, E_n \Rightarrow E$ and be

$$A(\theta) = 0; B(\theta) = 0; \dots$$

their associated conditions respectively where $A(\theta); B(\theta) \dots$ are polynomials not multiple of any complex polynomial. Then $E(\theta)$ is multiple of $\text{m.c.s.}[A(\theta), B(\theta), \dots]$

Proof. Factorize $\text{m.c.s.}[A(\theta), B(\theta), \dots]$ as product of un-factorizable polynomials

$$X(\theta); X_1(\theta); X_2(\theta); \dots$$

This passage is possible thanks to the unique factorization of polynomials. For every $X_n(\theta) \in \{X(\theta), X_1(\theta), X_2(\theta), \dots\}$ it is true that if $X_n(\theta) = 0$ then at least one among

$$A(\theta); B(\theta); \dots$$

Has value zero because $X_n(\theta)$ divides it. Moreover, from the hypothesis, if one among $A(\theta); B(\theta); \dots$ is equal to zero then also $E(\theta) = 0$ for the same values of $a, b, c, \dots (= \theta)$.

This means $X_n(\theta) = 0$ implies $E(\theta) = 0$.

Then apply theorem 2.9 on the polynomials $X_n(\theta)$ and $E(\theta)$, which satisfy the conditions requested, and obtain that $X_n(\theta)$ divides $E(\theta)$. So each of $X(\theta); Y(\theta); Z(\theta); \dots$

Divide $E(\theta)$ and as they are the factorization of $\text{m.c.s.}[A(\theta), B(\theta), \dots]$ if all of them divide $E(\theta)$ then also $\text{m.c.s.}\{A(a, b, c, \dots), B(a, b, c, \dots), \dots\}$ divides $E(a, b, c, \dots)$ $\square$

Q.E.D.

Remark 3.2. Now it is possible to answer the question:

Which feature do all the predicates σ which are true about rectangular triangles have in common?

Consider the generic triangle $A(0, 0)B(b, a)C(c, 0)$:

it is right if 1) $AB \perp BC$ or 2) $AB \perp CA$ or 3) $BC \perp CA$

1, 2, 3 are predicates and their associated conditions are

$$1) AB \perp BC : a^2 + b^2 - cb = 0$$

$$2) AB \perp CA : b = 0$$

3) $BC \perp CA : b = c$. the fact that these predicates σ are true about rectangular triangles means that the predicates "the triangle is rectangular on..." imply every predicate of σ . This let us apply the Autarky theorem

through which we know that the polynomials associated with all the true predicates σ have to be multiple of

$$(a^2 + b^2 - cb)(b - c)b$$

However the result above is not valid for predicates like "given a triangle ABC, if M is the middle point of BC then $AM = BC/2$ " because it is valid only if $AB \perp AC$ and not if $BC \perp CA$ or $AB \perp BC$.

4. AN EXAMPLE OF USAGE

The theorem offers a way to prove predicates on the base of the analysis of some particular cases. This way avoids the length of calculations required by the analytical demonstration and does not require the creativity which characterize the synthetic geometry, but it has little practical use because the nowadays computational programs can support an incredible computation complexity. Any demonstration made by this technique is composed of some passage as in the example that follows.

Theorem 4.1 (Existence of Euler's line). *In a general triangle orthocenter, baricenter and circumcenter are collinear.*

Proof. **Step 1**

If the triangle is isosceles, the demonstration is simple (all the three points belong to the axis of symmetry).

Step 2

We are going to write the problem on the cartesian plan to evaluate the maximum degree of the associated condition. We will use as generic triangle the one of vertexes $A(0,0)B(b,a)C(c,0)$ as in proposition 1.4. Find the coordinates of the three points depending on (a,b,c) :

Baricenter $G(\frac{b+c}{3}; \frac{a}{3})$

Orthocenter $H(b; \frac{b(c-b)}{a})$

Because it is the intersection of the height from $B : x = b$ and the one from $A : y = \frac{c-b}{a}x$

Circumcenter $C(\frac{c}{2}; \frac{(b^2 - cb + a^2)}{2a})$

Because it is the intersection of the axis of $AC : x = \frac{c}{2}$ and the one of $AB : y = \frac{(b^2 - xb + a^2)}{2a}$ (the last line is obtained as set of points whose distances from A and B are equal)

Step 3

In order to complete the proof we should apply the formula of three point collinearity and obtain an equation which is the associated condition of the thesis. Once shown that we obtain an identity the theorem is proved for every (a, b, c) . The condition is:

$$(x_H - x_G)(y_C - y_G) = (y_H - y_G)(x_C - x_G)$$

(Where x_H is the x-coordinate of H, y_H is the y-coordinate of H and the same for G and C). We have to evaluate the maximum possible degree of this equation by substituting the degree of the polynomials for the polynomials themselves.

(For example $y_C = \frac{(b^2 - cb + a^2)}{2a} = \frac{two}{one}$)

$$(x_H - x_G)(y_C - y_G) = (y_H - y_G)(x_C - x_G)$$

corresponds to

$$one(\frac{two}{one} - one) = one(\frac{two}{one} - one)$$

(The difference between two polynomials has a degree which is lower or equal to the one of the major of them and the product of two polynomials has degree which is lower or equal to the sum of their degrees)

$$\frac{three - three}{one} = \frac{three - three}{one}$$

$$four = four$$

The degree of the associated condition is lower or equal than four.

Reductio ad absurdum

At this point, if the associated condition is not an identity, it has degree which is minor or equal than 4 but, thanks to step 1, it is possible to apply Autarky theorem which says that the associated polynomial can be written as $P(a, b, c) = 0$ where $P(a, b, c)$ is multiple of the m.c.s. of the polynomials associated with the predicates: "ABC is isosceles on base AB", "ABC is isosceles on base CB", "ABC is isosceles on base AC"

If ABC is isosceles on AB, $AC=CB$, so $a^2 + b^2 - 2bc = 0$

If ABC is isosceles on CB, $AB=AC$, so $a^2 + b^2 - c^2 = 0$

If ABC is isosceles on AC, $AB=CB$, so $2b - c = 0$

(the three polynomials $(a^2 + b^2 - 2bc)$; $(a^2 + b^2 - c^2)$; $(2b - c)$ are real and irreducible so they are truly the C.A. with these predicates respectively and they respect the conditions required to apply the Autarky Theorem).

This means that $P(a, b, c)$ is multiple of
m.c.s. $\{(a^2 + b^2 - 2bc); (a^2 + b^2 - c^2); (2b - c)\} =$

$$(a^2 + b^2 - 2bc)(a^2 + b^2 - c^2)(2b - c)$$

But this is absurd because a polynomial of degree four cannot be multiple of one of degree five. So the associated condition must be an identity and the thesis is proved.

Q.E.D.

The demonstration schema is:

Find as many particular cases such that their predicate E imply the truth of the thesis as possible

Write the C.A. of each of these predicates and evaluate the degree D of their m.c.s.

Evaluate the thegree d of the C.A. with the thesis

If $D > d$ the thesis is proved. $\square$

5. THE UNIVERSAL TRIANGLE

One important application of the definitions given in the first part of this article is the creation of an "universal triangle", id est a triangle which does not own any property which is not common to all the triangles. This creation is possible only through the application of this result of high abstract algebra:

Theorem 5.1. (*LindemannWeierstrass*) *Let $a_1, \dots, a_n$ be algebraic numbers which are linearly independent over the rational numbers Q , then $e^{a_1}, \dots, e^{a_n}$ are algebraically independent over Q .*

As the set A of the algebraic numbers is the algebraic clousure set of Q , this theorem can be extended to all the equations with coefficients A , so that $e^{a_1}, \dots, e^{a_n}$ are algebraically independent over A .

Let ABC be the triangle such that $A(0, 0)$ $B(e, e^{\sqrt{2}})$ $C(e^{\sqrt{3}}, 0)$. Thanks to Lindemann-Weierstrass theorem, there is no polynomial equation with coefficients in A such that $e, e^{\sqrt{2}}, e^{\sqrt{3}}$ belongs to its solutions. As said above the fact that all the construction formula in the cartesian plane (perpendicular and parallel lines, middle points, bisectors, incircles,...) are reducible to polynomials (do not contain trascendental function) implies that the associated condition with a predicate of syntetic geometry is always a polynomial equation with algebrical coefficients.

This means that the only predicates which are true about this universal triangle are identities, so they are generally true (the only condition is that the predicate which we are dealing with can be expressed in terms of the standard geometric constructions). Every predicate (which depends on three points) whose C.A. can be amenable to a polynomial equation with algebraic coefficients is true if, and only if, it is true about the universal triangle: every predicate of syntetic geometry can be proven just for the universal triangle to demonstrate its validity overall. As it is possible to find an arbitrarily long list of algebraically independent numbers

$$e, e^{\sqrt{2}}, e^{\sqrt{3}}, e^{\sqrt{5}}, e^{\sqrt{7}}, e^{\sqrt{11}} \dots$$

it is possible to find an arbitrarily complexed figure, which depends on an arbitrarily big set of variables, which has the same features of universality of the universal triangle.

6. INHOMOGENEITIES

Definition 6.1. A function $f(x, y, z, \dots)$ is called *homogeneous* of degree a if, for every k :

$$f(kx, ky, kz, \dots) = k^a f(x, y, z, \dots)$$

How can we know that the C.A. with a predicate E is an homogeneous polynomial equation?

The analytical plane needs the possibility to introduce an arbitrary "unit of measure" u such that the validity of the geometry theorems does not depend on the choosing of u . This is made possible by the fact that, being h the distance between two points, if we apply a similar transformation, which means if we multiply for the same quantity k_0 all the coordinates of the points of a geometric figure, the value of h is multiplied for k_0 . This means that if a geometric figure is built on the points $A, B, C, D \dots$ whose coordinates depend on the set of variables $S\{a, b, c, \dots\}$, and $h(a, b, c, \dots)$ represents the length of a segment, then $h(k_0a, k_0b, k_0c, \dots) = k_0h(a, b, c, \dots)$.

All the quantities which express a distance are degree one homogeneous functions (note that the variables that we have used so far, a, b, c can be considered degree one homogeneous polynomials and they represent the distance of some points from the axes) while all the quantities which

represent an angular coefficient should be homogeneous of degree zero, because an angular coefficient is a quotient of two lengths. This means that whenever a predicate expresses an equality between two lengths or two angular coefficients, its associated condition is an homogeneous polynomial equation (call Homogeneous equation the equality between two homogeneous function of the same degree). This does not mean that an equality between lengths has degree one:

Example 6.2. The function $\frac{a^2}{b}$ is homogeneous of degree 1, so it can represent the length of the segment AB . Being $AC = c$, the equation $AC = AB$ corresponds to $a^2 = bc$ which is homogeneous of degree two.

Once we have determined that an equation is homogeneous, we can apply some properties of the homogeneous functions. It has already been proven that:

- 1) the product of two homogeneous functions is still an homogeneous function
- 2) the quotient of two homogeneous functions is still an homogeneous function

It is useful to add one more property whose demonstration is more difficult:

Lemma 6.3. *Let $f(x, y, z, \dots) = f(\theta)$ an homogeneous polynomial. Let $f(\theta) = P(\theta)Q(\theta)$ where P, Q are polynomials. Then $f(\theta)$ is the product of two homogeneous polynomials.*

Proof. Let n, g, h be the degrees of $f(\theta), P(\theta), Q(\theta)$ respectively. Since the product of two polynomials has the degree of the sum of their degrees, $n = g + h$. Let $P(\theta) = O(\theta) + o(\theta)$ and $Q(\theta) = M(\theta) + m(\theta)$ where $O(\theta)$ is the part of $P(\theta)$ made by the monomials of degree g , and $M(\theta)$ is the part of $Q(\theta)$ made by the monomials of degree h . Note that $O(\theta), M(\theta)$ are homogeneous of degrees g, h respectively. This process can be repeated on $o(\theta), m(\theta)$ so that $o(\theta) = o_1(\theta) + o_2(\theta) \dots$ where $\forall \mu < g, o_\mu(\theta)$ is an homogeneous polynomial of degree $n - \mu$ (the same is valid for $m(\theta) = m_1(\theta) + m_2(\theta) + \dots$).

Now, from the definition of homogeneity,

$$\forall k, \forall \theta, P(k\theta) = k^g O(\theta) + k^{g-1} o_1(\theta) \dots$$

$$\forall k, \forall \theta, Q(k\theta) = k^h M(\theta) + k^{h-1} m_1(\theta) \dots$$

These can be considered polynomials in the only variable k by fixing the quantity θ .

$$f(\theta) = P(\theta)Q(\theta) = O(\theta)M(\theta) + o_1(\theta)M(\theta) + m_1(\theta)O(\theta)...$$

So

$$f(k\theta) = O(k\theta)M(k\theta) + o_1(k\theta)M(k\theta) + m_1(k\theta)O(k\theta)...$$

$$k^n f(\theta) = k^n O(\theta)M(\theta) + k^{n-1} o_1(\theta)M(\theta) + m_1(\theta)O(\theta)...$$

Note that, in the last equation, all the terms except for $k^n O(\theta)M(\theta)$, have a degree less than n with respect to the variable k . So, the only way to equalize these one-variable polynomials is:

$$\forall \theta, f(\theta) = O(\theta)M(\theta)$$

where $O(\theta), M(\theta)$ are homogeneous. □

This proof by itself does not argue that every homogeneous polynomial is dividable only for homogeneous polynomials, however it is simple to prove the latter theorem with this result.

Theorem 6.4. *Let $f(\theta)$ be an homogeneous polynomial and let $G(\theta)$ so that $G(\theta)$ divides $f(\theta)$. Then $G(\theta)$ is homogeneous.*

Proof. Let $f(\theta) = P(\theta)Q(\theta)$. For lemma 6.3 $f(\theta) = M(\theta), O(\theta)$ are homogeneous. Repeat this process on $M(\theta)$ and $O(\theta)$ we obtain un-factorizable polynomials. This algorithm continues until we obtain the factorization of $f(\theta)$ in un-factorizable polynomials. As the algorithm produces only homogeneous polynomials, $f(\theta)$ can be factorized into a product of homogeneous polynomials.

$$f(\theta) = f_1(\theta)f_2(\theta)f_3(\theta)...$$

Let $R(\theta)$ be an inhomogeneous polynomial which divides $f(\theta)$. $R(\theta)$ should be multiple of some $f_n(\theta)$ and no other polynomials but this is absurd because the product of homogeneous polynomials must be an homogeneous polynomial. □

The importance of this result resides in the fact that now if we know that the C.A. with a predicate E is an homogeneous polynomial equation $P(a, b, c, \dots) = 0$, if we prove that $\forall a, b, c, \dots$

$$G(a, b, c, \dots) = 0 \Rightarrow P(a, b, c, \dots) = 0$$

where $G(a, b, c, \dots)$ is an *inhomogeneous* polynomial which respects all the

conditions to apply the theorem 2.9, then E is established. That is because the theorem 2.9 states that $G(a, b, c, \dots)$ divides $P(a, b, c, \dots)$, but for theorem 6.4 this is contradictory unless $P(a, b, c, \dots) = 0$ is an identity.

Proposition 6.5. *Let E be a predicate on the variables $a, b, c, \dots = \theta$. Let $C.A.$ with E be an homogeneous polynomial equation. If E is valid in the case $a^2 = b$, then E holds always.*

This is clearly because $a^2 - b$ is not an homogeneous polynomial. This is an example of substitution which can be made to simplify the calculus of an analytic geometry problem. When dealing with a geometric predicate which depends on the variables $a, b, c, d, \dots$ we can always make a substitution $P(a, b, c, d, \dots) = G(a, b, c, d, \dots)$ (inhomogeneous polynomial equation) to simplify the calculus. In fact if a predicate E can be proved using the substitution S , then $S \Rightarrow E$ and this form let us apply the theorems in the previous sections.

REFERENCES

- [B] F. Botaccin, *Introduzione alla geometria algebrica*, Universita dagli Studi di Padova, Padova, 2007.
- [R] L. Rigolli, *Hilbert Nullstellensatz e alcune sue conseguenze geometriche*, Graduation thesis, Universita degli Studi di Bologna, Bologna, 2011.
- [W] Wikipedia, *Teorema di Lindemann-Weierstrass*, last visit 12/28/16.
- [W] Wikipedia, *Numero algebrico*, last visit 12/28/16.

E-mail address: `davide.maran@mail.polimi.it`